

Hidden Local Field Theory and Dileptons in Relativistic Heavy Ion Collisions

Gerald E. Brown,¹ Masayasu Harada,² Jeremy W. Holt,^{1,3} Mannque Rho,^{4,5} and Chihiro Sasaki³

¹*Department of Physics and Astronomy, State University of New York, Stony Brook, NY 11794, USA*

²*Department of Physics, Nagoya University, Nagoya, 464-8602, Japan*

³*Physik-Department, Technische Universität München, D-85747 Garching, Germany*

⁴*Institut de Physique Théorique, CEA Saclay, 91191 Gif-sur-Yvette cedex, France*

⁵*Department of Physics, Hanyang University, Seoul 133-791, Korea*

(Dated: January 12, 2009)

The notion of “hadronic freedom” is introduced based on the vector manifestation of hidden local symmetry and is used to suggest that the dileptons measured in relativistic heavy-ion collisions do not provide *direct* information on the spontaneous breaking of chiral symmetry and hence on the mechanism for mass generation of light-quark hadrons. We give arguments how the dileptons emitted from those vector mesons whose masses are shifted à la Brown-Rho (BR) scaling by the vacuum change in temperature – as in heavy-ion collisions – and/or density – as in cold compressed matter – could be strongly suppressed in the hadronic free region between the chiral restoration point and the “flash point” at which the vector mesons recover $\gtrsim 90\%$ of their free-space on-shell masses and the full strong coupling strength. It may seem ironical that the very mechanism i.e., the vector manifestation in hidden local symmetry, that is to make the mass drop and modify the spectral function is in turn responsible for the dilepton suppression. A possible falsification of this drastic prediction will be indicated. We also briefly discuss the potential role played by the holographic dimension intrinsic in gravity-gauge duality that provides a unified field theory description of hadrons – both mesons and baryons – under extreme conditions. Baryons arise as coherent states of pions and vector mesons of holographic QCD which define the ground state or the “vacuum” on which mesonic excitations could undergo BR scaling.

I. INTRODUCTION

In QCD, most of the mass of the ground-state light-quark hadrons, e.g., $\gtrsim 95\%$ of the ρ meson mass, is understood to be generated dynamically by the spontaneous breaking of chiral symmetry in the complex vacuum, so it is natural to think that by “cleansing” the vacuum, the masses could disappear as the broken symmetry is restored. Based on this general consideration, a prediction was made in 1991 that the masses of light-quark hadrons would drop proportionally to some power of the quark condensate at high temperature and density [1]. This in-medium property of hadrons is referred to in the literature as “Brown-Rho (BR) scaling.” It was thought that the “unbreaking” of the symmetry, or equivalently the “shedding” of the mass, can be done by heating the hadronic system to a high temperature (and/or by compressing the matter to a high density). To unravel the vacuum change induced specifically by temperature, efforts have been made to look at the properties of the vector mesons, i.e. ρ and ω , in heavy ion collisions where the temperature could be raised to hundreds of MeV at which chiral restoration is predicted by QCD to take place. It was thought that the properties of the ρ (and ω) meson inside strongly interacting media could be effectively probed with dilepton pairs, which couple to the vector mesons and provide, free of final state interactions, a snap-shot of the vector meson propagating in the medium. Due to the mass shedding in a hot medium,

one naturally expected to observe a downward shift of the invariant mass of the vector meson in the spectral functions. The measurement of dileptons, which is the classical tool for discovering new resonances in particle physics, is used to see how the known ρ resonance in free space moves in a hot medium.

Several such experiments have been performed recently. For reasons which will become clear below, we will focus specifically on the NA60 [2] experiment, although our discussion applies to other dilepton measurements such as CERES, etc. What came out of the experiments so far performed is that no vector mesons affected by the vacuum change due to temperature and density as predicted by BR scaling are unequivocally “seen” by the dileptons.

In this paper, using a set of arguments, some established and some conjectural, based on the structure of hidden local symmetry in low-energy hadronic interactions, we develop the scenario that dileptons emitted from the vector mesons whose properties are BR-scaled are strongly, if not completely, suppressed and the dileptons observed in heavy-ion experiments reflect on-shell behavior of the vector mesons only modified by the mundane nuclear many-body interactions [3] but not by the change of the vacuum due to temperature and/or density. This means that if our scenario is correct, *the BR-scaled vector mesons are almost, if not entirely, “invisible” to the virtual photon.* As briefly commented in the conclusion section, a similar consideration – with a small variation – could apply to dileptons produced in such cold

dense environments as expected to be provided at the future FAIR/GSI facility.

We propose as a promising tool to unravel the chiral symmetry structure of hadronic matter a holographic hidden local symmetry framework in which the holographic dimension allows, with a single 5D hidden local symmetry action, a unified description in medium of both mesonic and baryonic excitations treated on the same footing. Baryons emerge here as coherent states of pions and vector mesons which define the ground state or the modified “vacuum” on which fluctuating vector mesons undergo BR scaling in hot and/or dense medium. A consistent treatment along this line, yet to be made, could lead to a totally different picture from that currently accepted of hot and dense baryonic matter.

II. THE NOTION OF HADRONIC FREEDOM

The key proposition in our work is that in a certain range of temperature (and density [4]) to be specified below between the critical T_c at which a chiral transition takes place and what we will call “flash temperature” $T_f < T_c$, *all* hadron interactions become weak, as a consequence of which vector-meson coupling to the virtual photon γ^* is suppressed. This is associated, in a manner to be precisely specified below, with the “vector manifestation” (VM) phenomenon in hidden local symmetry (HLS) developed by Harada and Yamawaki [5]. We first briefly summarize the predictions of HLS as one approaches T_c from below that is relevant to the problem at hand and then introduce as clearly as possible the notion of “hadronic freedom” that plays the principal role in our discussions.

A. Hidden local symmetry in low-energy strong interactions

We start with a brief sketch of the 5 dimensional hidden local symmetry action that presumably encodes all low-energy hadron dynamics involving integer spin (0, 1) and half-integer (1/2, 3/2, ...) spin excitations treated on the same footing. One can arrive at an action of the same structure both bottom-up and top-down.

The well-established theory for strong interactions at low energy is chiral perturbation theory for the (pseudo-) Nambu-Goldstone pions $U = e^{2i\pi/F_\pi}$, the effective Lagrangian of which has the form

$$\mathcal{L}_{eff} = \frac{F_\pi^2}{4} \text{Tr}(\partial_\mu U \partial^\mu U^\dagger) + \dots \quad (1)$$

where the ellipsis stands for higher derivative and quark mass terms. At energies much less than the vector-meson excitation $m_V \sim 800$ MeV, when treated in a systematic expansion, this represents QCD [6]. The validity of this theory is however highly limited to low energy, so if one wants to study the vector mesons whose masses are much

higher than that of the pion, chiral perturbation theory with pions alone loses its power. The question is how to extend the model-independent approach to higher-energy scale. We will be particularly concerned with the situation where the vector-meson mass decreases from its free-space mass m_V to one near the pion mass which is zero in the chiral limit [7]. In our approach, this occurs at the chiral phase transition. This means that we will be dealing with the possibility – which will be realized in our model – that the vector meson mass becomes comparable to the pion mass and hence both the vector meson and the pion need to be treated on the same footing. This can be most readily done if one exploits local gauge invariance [5, 8, 9]. Seen from the holographic point of view, this is natural as we will argue below.

The astute observation for our development is that the chiral field [10]

$$U = e^{\frac{2i\pi}{F_\pi}} \in G/H \quad (2)$$

where π denotes the Nambu-Goldstone (NG) bosons associated with the spontaneous breaking of chiral G symmetry to H with $G = SU(N_f)_L \times SU(N_f)_R$ and $H = SU(N_f)_{V=L+R}$, has a redundancy when written in terms of the coordinates for the L and R symmetries,

$$U = e^{\frac{2i\pi}{F_\pi}} = \xi_L^\dagger \xi_R \quad (3)$$

with

$$\xi_{L,R} = e^{i\sigma/F_\sigma} e^{\mp i\pi/F_\pi}, \quad [\pi = \pi^a \tau_a/2, \sigma = \sigma^a \tau_a/2] \quad (4)$$

The redundancy can be elevated to a local gauge invariance by noting that the chiral Lagrangian built with U is invariant under the local transformation

$$\xi_{L,R} \rightarrow h(x) \xi_{L,R} \quad (5)$$

with $h(x) \in SU(N_f)_V$ and introducing a gauge field $\rho_\mu = \rho_\mu^a \tau^a/2$ that transforms

$$\rho_\mu \rightarrow h(x)(\rho_\mu - ig\partial_\mu)h^\dagger(x). \quad (6)$$

The field σ represents the redundancy in the low-energy theory. Now when written in the lowest order in derivatives in terms of the covariant derivatives $D_\mu \xi_{L,R} = (\partial_\mu - ig\rho_\mu)\xi_{L,R}$, the resulting local gauge invariant Lagrangian $\mathcal{L}'[D_\mu \xi_{L,R}]$ is identical to the chiral Lagrangian (1) when gauge fixed to unitary gauge corresponding to $\sigma = 0$. One says that $\mathcal{L}'$ is “gauge-equivalent” to $\mathcal{L}_{eff}$. To elevate the energy scale from that of the current algebra scale to that of the vector mesons, one adds the kinetic energy term to make the field propagate,

$$\mathcal{L}_{HLS} = \mathcal{L}'[D_\mu \xi_{L,R}] - \frac{1}{2} \text{Tr}(\rho_{\mu\nu} \rho^{\mu\nu}) + \dots \quad (7)$$

The σ can be identified with the Nambu-Goldstone (NG) scalars that emerge from the spontaneous breaking of the H symmetry and that are higgsed to give masses to the

vector mesons. This is hidden local symmetry theory proposed in [11, 12] and later quantized [5] that we shall refer to as HLS₁.

For our development, it is important that the notion of hidden local symmetry can be extended to an infinite number of hidden local fields. We suggest that HLS₁ is a truncated version of a hidden local symmetry theory with an infinite tower of vector fields. As such, HLS₁ can be viewed as a realistic model of QCD.

It is clear from the construction of HLS₁ that there is nothing that prevents adding an infinite number of redundancies and construct a theory with an infinite number of gauge fields. There is of course the problem of non-uniqueness since the construction would lead to a large number of possibilities. One such construction which resembles the nearest neighbor interaction in condensed matter physics is the construction based on “theory-space locality” [9] on lattice which in the continuum limit leads to a five-dimensional (5D) Yang-Mills action defined in a curved space [13]

$$S_{YM} = \int d^4x dz \sqrt{g} \frac{1}{2g_5(z)^2} \text{Tr}(F_{AB}F^{AB}) + \dots \quad (8)$$

with $A = 0, 1, 2, 3, z$. The ellipsis stands for higher derivative terms. Implemented with the Chern-Simons action S_{CS} that encodes anomalies, this may be considered as a “dimensionally deconstructed” QCD [13] emerging bottom-up from the current algebra Lagrangian. Note that this action contains not only the Goldstone bosons and spin-1 excitations but also baryons as instantons, i.e., coherent states of the Goldstone boson and vector mesons, all on the same footing.

What is most intriguing is that a 5D YM action plus the Chern-Simons identical in form to the action (8) arises also top-down from string theory. Indeed an action of the form (8) was recently constructed by Sakai and Sugimoto [14] using D4/D8-D $\bar{8}$ branes valid in the limit that $N_c \rightarrow \infty$, $\lambda \equiv g_{YM}^2 N_c \rightarrow \infty$ and in the chiral limit. Now reduced to four dimensions (4D) by the usual Kaluza-Klein procedure, the action can be phrased in terms of an infinite tower of vector and axial vector mesons describing $U(N_f)_L \times U(N_f)_R$ chiral symmetry which may be rewritten in a local gauge invariant form. We shall refer to the infinite-tower local symmetry theory as HLS _{∞} . Various approximations enter in arriving at the 5D action such as for instance the “probe approximation” $N_f/N_c \ll 1$. But what is robust and generic is that low-energy hadron dynamics can be encapsulated in a hidden local symmetric action with an infinite tower of gauge fields. Notice however that Eq.(8) bottom-up is defined in the gauge sector whereas top-down is in the gravity sector. Generating a gauge invariant theory from a non-gauge theory as in the bottom-up approach does not lead to a unique theory and hence can be totally arbitrary unless constraints are imposed. On the other hand, the top-down approach is well-defined in the given limit but its UV completion is to a certain part of string theory which may have nothing to do with QCD.

Furthermore, while the problem we are interested in involves properties that are intricately locked to the chiral order parameter, i.e., quark condensate, which changes with temperature and density, the large N_c and λ limits suppress the temperature and/or density dependence in the physical variables. One would have to calculate terms higher order in $1/N_c$. But that would require calculating string loop corrections in the gravity sector which is at present infeasible. For this reason, we are unable to look at vector meson masses as temperature increases since in the large N_c limit, the masses do not move. Nonetheless, as we will see below, the infinite tower structure has an extremely important implication on vector dominance in hadron electromagnetic form factors and this feature is expected to play a crucial role in understanding dilepton production process in heavy ion collisions.

Let us, for the sake of arguments, imagine that we have an HLS _{∞} in which all pertinent $1/N_c$ and $1/\lambda$ corrections can be included and which provides a realistic description of low-energy QCD. Assume further that as temperature is increased, only the lowest vector meson $\rho_0 = \rho(770)$ drops in mass while the higher lying vector mesons remain more or less unmodified. This is most likely very difficult to confirm, but it is plausible. In this case, we could imagine formally integrating out *all* vector mesons except ρ_0 and write the resulting action as a suitable power series in terms of covariant derivatives of the ρ_0 field to get a hidden local symmetric effective action. How to do this consistently has been proposed by Harada, Matsuzaki and Yamawaki [15]. What results would be identical in form to the HLS₁ theory of [5, 11, 12]. This would justify HLS₁ as a potentially realistic model for low-energy dynamics in high temperature, particularly if the ρ_0 mass does come down to that comparable to the pion mass which we will argue is what happens at high temperature in heavy ion collisions. As suggested in [5], the local gauge symmetry plays a crucial role in treating this situation.

B. The vector manifestation (VM)

In free space at $T = 0$, the vector meson mass m_{ρ_0} can be taken to be very heavy compared to the pion mass, i.e., $m_{\rho_0}/m_\pi \gg 1$, particularly in the chiral limit. In this case, there are various different ways of introducing vector mesons – with or without local gauge symmetry – which are all equivalent at tree order. The hidden local symmetry approach HLS₁ on the contrary takes the vector meson mass to be of the same order as the pion mass, that is, in the chiral power counting,

$$m_\rho \sim m_\pi \sim \mathcal{O}(p) \quad (9)$$

which is equivalent to taking the gauge coupling g going as

$$g \sim \mathcal{O}(p). \quad (10)$$

In the framework adopted here where the ρ mass does decrease, it is very likely that this power counting holds, rendering chiral perturbation expansion justified in the vicinity of the chiral transition point T_c where the vector meson mass goes to zero. But it is not obvious why it should hold in free space where m_ρ/m_π is not of order 1. Somewhat surprisingly, however, it does turn out that chiral perturbation calculation including the vector meson works fairly well in free space as shown in [5]. This indicates that the chiral perturbation which is justified in large N_c describes the realistic $N_c = 3$ world as an extrapolation.

Given the systematic counting rule, one can then develop chiral perturbation. This was done to one-loop order [16]. In the parameter space of g , F_π and $a = (F_\sigma/F_\pi)^2$ (quark masses should of course be additional parameters if we go beyond the chiral limit.), the renormalization group analysis of HLS₁ theory to one loop order reveals a variety of fixed points. This reflects the well-known non-uniqueness of gauge symmetric theories constructed from non-gauge theories [17]: local gauge symmetric theories can flow, unconstrained, to a variety of different fixed points. In order to identify the flow to QCD among them, Harada and Yamawaki match à la Wilson the vector and axial-vector correlators of HLS₁ to those of QCD at a matching scale Λ_M . The matching then determines the HLS₁ parameters (g, a, F_π) in terms of the QCD variables, namely, α_c , N_c , $\langle \bar{q}q \rangle$, $\langle G_{\mu\nu}^2 \rangle$ etc. that figure in the OPE of the correlators.

Now when one imposes the condition that at the chiral phase transition characterized by $\langle \bar{q}q \rangle = 0$, the vector correlator Π_V equal the axial-vector correlator Π_A – which is what one expects in QCD, one can identify the fixed point (denoted by bar) that corresponds to QCD, i.e.,

$$(\bar{g}, \bar{a}) = (0, 1). \quad (11)$$

This point is called the “vector manifestation” fixed point [5]. Although this fixed point is derived at one-loop order, it can be seen with a little effort that it is valid to all orders. The parametric pion decay constant F_π does not figure in the fixed point (11) because there is nothing special about the flow of F_π , so at the fixed point, it does not have any special role as it remains non-vanishing. However the on-shell (physical) pion decay constant f_π which receives loop corrections Δ below the on-shell ρ mass,

$$f_\pi = F_\pi + \Delta \quad (12)$$

should go to zero at T_c because it must track the quark condensate. Note that this must occur by cancellation. What is important is that as T goes to T_c [18] (and as density n goes to the critical density n_c [19]), the hadronic matter flows to the VM fixed point (11),

$$\begin{aligned} g^* &\sim \langle \bar{q}q \rangle^* \rightarrow 0, \\ (a^* - 1) &\sim \langle \bar{q}q \rangle^{*2} \rightarrow 0. \end{aligned} \quad (13)$$

Here and in what follows the asterisk stands for temperature (or density) dependence. This means that the parameters of the Lagrangian have an intrinsic dependence on temperature locked to the property of the condensate $\langle \bar{q}q \rangle$. This has the key consequence that the *parametric mass* of the vector meson very near the VM fixed point behaves as [20]

$$M_\rho^{*2} = a^* F_\pi^{*2} g^{*2} \sim \langle \bar{q}q \rangle^{*2}, \quad (14)$$

so will go to zero. This is the *statement of BR scaling* sharpened by HLS₁.

Now what about the physical (or pole) mass of the vector meson m_ρ in medium? For this one has to compute thermal loop terms which will necessarily involve the gauge coupling, so

$$m_\rho^{*2} = a^* g^{*2} F_\pi^{*2} + g^{*2} B \quad (15)$$

where B is a smooth function of temperature given by thermal loop corrections. As $g^* \rightarrow 0$, both terms go to zero as $\sim g^{*2}$, so that BR scaling should still hold for the physical mass. It is however to be noted that far away from the VM fixed point, there is nothing that suggests that the physical mass must be dropping according to BR scaling. In fact, it can go up or down or even remain more or less unchanged depending on the thermal loop corrections involving complicated many-body nuclear dynamics. In order to “see” BR scaling in terms of HLS₁ in temperatures away from the VM fixed point, therefore, one needs to bring in baryons via topology in addition to the bosonic degrees of freedom that are present explicitly. Such a calculation remains to be done. The same consideration holds in cold dense matter as discussed in the conclusion section.

C. The hadronic freedom

In the close vicinity of the VM fixed point, the interactions are likely to be weak since the gauge coupling goes to zero. The main reason for this is that in HLS₁, the strongest interaction at low energy which is in the p-wave channel, i.e., the ρ channel, is suppressed by the vanishing g . Further, the s-wave $\pi\pi$ interactions are governed by derivative interactions (in the chiral limit), so should be weak at low energy. There may be other degrees of freedom not explicitly taken into account in HLS₁ – such as quasiquarks or possible scalars – but the quasiquark interactions mediated by the vector mesons are also suppressed by the gauge coupling and quasiquark-pion interactions are derivative-coupled, so suppressed by the power counting. Thus it seems reasonable to take *all* hadronic interactions to be weak near the VM fixed point. Now how far down in temperature can one ignore the interactions? In principle, the intrinsic temperature dependence will be running as temperature is varied. For instance, the gauge coupling in the chiral limit will scale up from zero at T_c to near its free space value at some temperature which we shall call “flash temperature” T_f .

Flash temperature T_f

The flash temperature T_f can be defined following the work of Shuryak and Brown [21] of the STAR data [22]. Shuryak and Brown determined the flash temperature to be the temperature at which the ρ was 90% on-shell, namely, when the ρ mass is 700 MeV. Whereas the flash temperature is independent of the centrality, this is not the case for the freezeout temperature $T_{freezeout}$. In fact $T_{freezeout}$ decreases as the centrality increases. Thus, given $T_{freezeout} \simeq T_f$ for peripheral collisions, $T_{freezeout} < T_f$ for central collisions and during the time the system is between these two temperatures the pions from ρ and a_1 decay will be rescattered as it is no longer possible from their detection to work backward to the parent vector mesons.

It is possible to pin down the numerical value of the flash temperature from lattice calculations. We shall extract this quantity from Miller's lattice calculation [23, 24] which we analyzed in [25]. The result is

$$T_f \approx 120 \text{ MeV}. \quad (16)$$

How this result comes about can be summarized as follows. As pointed out in [25], in Miller's lattice calculation of the gluon condensate, the soft glue starts to melt at $T \approx 120$ MeV. The melting of the soft glue, which breaks scale invariance as well as chiral invariance *dynamically* and is responsible for Brown-Rho scaling, is completed by T_c at which the particles have gone massless according to the VM. One can see that the gluon condensate at $T \sim 1.4 T_c$ is as high as that at $T \sim T_c$. This represents the hard glue (or "epoxy") which breaks scale invariance *explicitly* but has no effect on the hadron mass. We see that the melting of the soft glue is roughly linear, implying that the meson masses drop linearly with temperature. [24]

How the (bare) parameters of the Lagrangian scale *intrinsically* in temperature (and/or density) is not quantitatively known except in the very close vicinity of the VM fixed point. In the absence of guidance, theoretical or experimental, we take, as the first step, the simplest scenario which is to ignore interactions *entirely* between the chiral restoration point T_c and the flash point T_f and let the relevant massless degrees of freedom coming down from above T_c [26] flow without interactions as the system cools to T_f . We then assume that hadrons go (nearly free-space) on-shell at T_f recovering their (nearly) full interaction strength and free-space mass. Corrections due to increasing temperature as one approaches T_f – involving both the intrinsic dependence and the thermal loops – need of course to be computed at the next stage of refinement. This region between T_c and T_f will be dubbed "hadronic freedom regime." [27]

We must admit that the notion of hadronic freedom is a strong assumption which needs to be confirmed. At present, there are no theoretical tools to check this notion in unambiguous ways. As pointed out below, it could be validated or refuted by experiments. All we can say at

the moment is that while there are indirect evidences for it, there are no "smoking gun" evidences against the notion of hadronic freedom either. What we propose to do in this article is to *posit* the hadronic freedom and then see how far we can go in confronting nature. We will then present our interpretation of its consequence on dileptons in heavy ion collisions. Our scenario, which is drastically different from the conventional scenario, has the merit of providing a possible, albeit indirect, link between the manifestation of BR scaling in low-energy nuclear structure physics – some of which will be mentioned in the conclusion section – and possibly direct manifestation or precursor effects in heavy ion processes.

D. Vector dominance and infinite tower

In considering dileptons in heavy ion collisions, it is customarily assumed that vector dominance holds in medium at *all* temperatures and densities. In HLS₁, vector dominance does hold in free space but not in hot/dense medium. In fact, vector dominance is violated at $T = T_c$, with a drastic effect on dileptons in hot/dense medium. We briefly review how this comes about and then discuss what happens in the presence of the infinite tower of vector mesons figuring in holographic QCD that follows from gravity/gauge duality in string theory. We believe that the infinite tower structure encoded in the holographic direction which treats both mesons (spin 0 and 1) and baryons on the same footing, when $1/N_c$ corrections are suitably handled, could ultimately clarify the behavior of hadrons in high temperature and/or density.




FIG. 1: Leading contributions to the electromagnetic form factor of the pion. (a) direct $\gamma\pi\pi$ and (b) $\gamma\pi\pi$ mediated by ρ -meson exchange. In terms of HLS₁ Lagrangian, (a) is proportional to $(1 - a/2)$ and (b) is proportional to $a/2$.

In HLS₁ theory, the photon couples to the charged pion as in Fig.1: (a) direct $\gamma\pi\pi$ with the coefficient $(1 - a/2)$ and (b) $\gamma\pi\pi$ mediated by ρ -meson exchange with the coefficient $a/2$. It turns out that $a = 2$ in medium-free ($T = n = 0$) space which gives the KSFR relation correctly. This value completely suppresses the direct photon coupling Fig.1(a), giving full strength to Fig.1(b). This is the famous lowest vector-meson dominance (LVD for short) of the pion form factor, i.e. ρ -meson dominance in Sakurai's sense [28]. Note however that $a = 2$ is not on the trajectories of the renormalization group equation for a , so it is in some sense an

accident [29]. Thus unsurprisingly, this LVD is violated – in fact “maximally” – as $T \rightarrow T_c$: As a approaches 1, the direct coupling enters with the same strength as the vector exchange term [18, 30]. This violation of LVD is argued below to be largely, if not wholly, responsible for the suppression of dileptons in the hadronic freedom regime.

An interesting possibility is that we might be able to understand the violation of LVD in HLS_1 in terms of the infinite tower in HLS_∞ . In the gravity/gauge dual model of Sakai and Sugimoto (SS) [14] which has the correct chiral symmetry structure of QCD, the pion form factor is vector dominated by the infinite tower:

$$F(q^2) = \sum_{n=0}^{\infty} \frac{g_{\rho_n} g_{\rho_n \pi \pi}}{m_{\rho_n}^2 - q^2} = \sum_{n=0}^{\infty} \xi_n \frac{g_{\rho_n \pi \pi}}{1 - q^2/m_{\rho_n}^2} \quad (17)$$

where

$$\xi_n = \frac{g_{\rho_n}}{m_{\rho_n}^2}, \quad (18)$$

and g_{ρ_n} and $g_{\rho_n \pi \pi}$ represent respectively the γ - ρ_n coupling and the ρ_n - $\pi\pi$ coupling. These couplings are determined in the large N_c and λ limit by the equations of motion for the ρ_n fields integrated over z (the fifth coordinate) which represents energy spread and are fixed once $N_c = 3$, f_π and the Kaluza-Klein (KK) mass M_{KK} are taken from the meson spectra.

At the photon point $q^2 = 0$, one has the charge sum rule for the charged pion,

$$F(0) = 1 = \sum_{n=0}^{\infty} \xi_n g_{\rho_n \pi \pi} = \bar{\xi} \sum_{n=0}^{\infty} g_{\rho_n \pi \pi} \quad (19)$$

where we have used the observation [14, 31] that

$$\xi_n = \frac{g_{\rho_n}}{m_{\rho_n}^2} = 0.272 \pm 0.001 \equiv \bar{\xi} \quad (20)$$

which holds for $n = 0, 1, 2, 3$. We have no proof but we will assume that it holds for *all* n . A surprising result obtained by Sakai and Sugimoto is that the sum rule (19) is saturated by the lowest four ρ 's, i.e., ρ_n with $n = 0, 1, 2, 3$,

$$F(0) = 1.31 - 0.35 + 0.05 - 0.01 = 1.00. \quad (21)$$

There is a tantalizing empirical evidence that this sum rule is satisfied in τ decay [32]. We will return to this observation later in connection with multi-pion-photon coupling.

It should be noticed that the formula in Eq.(17) is obtained by using the equations of motion for the vector meson fields. Thus the sum rule Eq.(21) holds only when the couplings have no appreciable momentum dependence. Furthermore, $1/N_c$ corrections which are expected to be important in the time-like region may not be negligible here. Actually, as shown in Fig. 2 of Ref. [33], the experimental data of the pion form factor are well fit

in the time-like region by the one obtained in the HLS_1 with only the $\rho_{n=0}$ contribution, in which $a \simeq 2$ is used and about 10% momentum dependence is added to g_ρ together with an appropriate $1/N_c$ correction to the width.

To understand the potential role of the higher members of the tower in the form factor, it is convenient to separate the form factor (17) into “low” and “high” components as

$$F(q^2) = F^{\text{low}}(q^2) + F^{\text{high}}(q^2) \quad (22)$$

with

$$F^{\text{low}}(q^2) = \sum_{n=0}^k \frac{g_{\rho_n} g_{\rho_n \pi \pi}}{m_{\rho_n}^2 - q^2}, \quad (23)$$

$$F^{\text{high}}(q^2) = \sum_{n=k+1}^{\infty} \frac{g_{\rho_n} g_{\rho_n \pi \pi}}{m_{\rho_n}^2 - q^2}. \quad (24)$$

The sum rule (21) suggests that we take $k = 3$. Now for low $|q^2| \lesssim m_{\rho(770)}^2$, we can drop q^2 in F^{high} (24) and find from (21)

$$F^{\text{high}}(q^2) \approx \sum_{n=3}^{\infty} \frac{g_{\rho_n} g_{\rho_n \pi \pi}}{m_{\rho_n}^2} \approx 0. \quad (25)$$

Our conjecture is that the statement $a \rightarrow 1$ as $T \rightarrow T_c$ is tantamount to the statement that in HLS_∞ the higher tower members for $n = 4, \dots, \infty$ in the large N_c limit and those for $n = 1, \dots, \infty$ with $1/N_c$ corrections included figure “maximally”. How the infinite tower figures in the nucleon EM form factor is illuminating in this connection. As discussed in [31], the VD in the nucleon form factor is violated in HLS_1 . In fact, it is described rather well by $a \sim 1$ in Fig.1 with the pion replaced by the nucleon field. In HLS_∞ , the higher members of the tower are found to play the role of the point-like coupling of HLS_1 [31].

Unfortunately the conjecture cannot, at present, be verified in the SS model. In HLS_1 , it is the vanishing of the quark condensate as $(a - 1) \propto \langle \bar{q}q \rangle^2 \rightarrow 0$ that “kills” the point coupling. However it is known that the quark condensate has no temperature dependence in the large N_c limit. This suggests that to verify the conjecture, $1/N_c$ corrections are needed: Although a part of the $1/N_c$ corrections could be included by hadronic loop corrections in the gauge sector [15], certain $1/N_c$ corrections in the gravity sector – so far inaccessible – may give important contributions.

III. SOME CONSEQUENCES THAT FOLLOW FROM THE HADRONIC FREEDOM

In this section, we address how our approach based on the hadronic freedom in HLS_1 fares with experiments in heavy ion collisions. Our attitude here will be as follows. We posit the hadronic freedom together with BR scaling that we infer from HLS_1 as the premise of our approach

and discuss the implications on a variety of processes that can be accessed by the approach. We will first treat the cases where our approach can make clear statements and then present what can be said about the dileptons in heavy ion collisions that have been controversial in the literature such as the NA60 data etc.

A. The STAR ρ^0/π^- ratio

As a case where the notion of the hadronic freedom makes a simple prediction, we recall how the STAR ρ^0/π^- ratio (STAR ratio for short) [34] can be understood.

In a recent experiment, STAR collaboration has reconstructed the ρ -mesons from the two-pion decay products in the Au + Au peripheral collisions at $\sqrt{s_{NN}} = 200$ GeV. The ratio found was

$$\frac{\rho^0}{\pi^-}|_{STAR} = 0.169 \pm 0.003(\text{stat}) \pm 0.037(\text{syst}), \quad (26)$$

almost as large as the $\rho^0/\pi^- = 0.183 \pm 0.001(\text{stat}) \pm 0.027(\text{syst})$ in proton-proton scattering. The near equality of these ratios was not expected, in that the ρ meson width $\Gamma \sim 150$ MeV in free space is the strongest meson re-scattering that one has. For instance, at SPS one needs about 10 generations in order to get the dileptons [35]. Furthermore, if one assumes equilibrium at the freeze-out, then the ratio is expected to come to several orders of magnitude smaller, say, $\frac{\rho^0}{\pi^-} \sim 4 \times 10^{-4}$ [36].

There may be a variety of explanations for this observation, some of which are along the standard line of thought from which our scenario appears to depart drastically. What distinguishes the result that follows from the hadronic freedom – which turns out to come out consistent with the experiment (26) – is that the mechanism is extremely simple if one assumes that the ρ^0 's are reconstructed at the flash point at which the ρ meson emerges from the hadronic freedom regime going through only one generation from T_c . We should however stress that no claim is made here that the proposed mechanism is either unique or the only viable explanation. It is just that our scenario has the merit to be simple and consistent within the proposed framework anchored on the hadronic freedom and the vector manifestation of HLS₁.

This matter was discussed in [37] to which we refer for details. Here we review it briefly for the sake of making our arguments self-contained.

The key point in our scenario is that because of the decreased width due to the hadronic freedom, the ρ meson is assumed to go through *only one generation* before it freezes out at the flash point in the peripheral collisions in STAR. There will be no equilibrium at the end of the first generation.

As the plasma decreases in temperature towards T_c from above, lattice gauge calculations[38] show a set of 32 $SU(4)$ degenerate vibrations[39]: scalar, pseudoscalar, vector and axial vector, i.e., the set of degrees of freedom

that figure in Weinberg's mended symmetry at the chiral transition point [40]. The widths become large as $T \rightarrow T_c$ from above. [41] We connect these with the $SU(4)$ group of mesons, including the ρ of the HLS₁ vector manifestation, becoming massless (except for a small spin-spin interaction effect) as T moves up to T_c from below. Aside from the background η and ω , these 32 mesons decay as they go on shell to give 65 pions (three of them being already present in the 32 $SU(4)$ mesons) [42, 43]. Thus, taking the 22 π^- mesons and subtracting 3, which are traced back to the vertex inside the plasma where the ρ^0 was formed, one has

$$\frac{\rho^0}{\pi^-} \approx \frac{3}{22 - 3} \approx 0.16 \quad (27)$$

in agreement with the experiment. As noted, this number is about the same as the ρ^0/π^- ratio measured in pp scattering. However, the vacuum ρ has a width of 150 MeV so its lifetime is only $\hbar/150 \simeq 4/3$ fm/c and in proceeding from T_c to the flash point where the ρ goes on shell, which we shall find to be at ~ 120 MeV, were the interactions unsuppressed, there would be several generations of $\rho \rightarrow 2\pi \rightarrow \rho$, whereas our estimates show that there are none; the ρ 's move from T_c to T_f without interaction. Thus had the ρ possessed its on-shell mass and width with $\sim$ five generations as in the standard scenario, the $\frac{\rho^0}{\pi^-}$ ratio would be closer to the much smaller equilibrium value.

B. Dense matter

There are certain consequences of the hadronic freedom that are more prominent in dense matter than in hot matter. Since dilepton experiments involve density in addition to high temperature, some of them need to be explained.

One of the most important applications of the notion of the hadronic freedom is to the role of kaon condensation in compact stars. A recent publication addresses this matter in conjunction with black-hole formation and a possible cosmological consequence [44].

While the phase structure of dense matter is not well understood, the first important phase change that can take place in dense matter above the nuclear matter density n_0 and that can be treated reliably thanks to the VM is kaon condensation at a density $\sim 3n_0$. This phase transition is considered to be responsible for the ultimate fate of compact stars, e.g., the formation of stable neutron stars or collapse into black holes [44]. It takes place because of the decrease of the kaon mass and the increase of the electron chemical potential at increasing matter density: Matter density drives the mass of the kaon downwards and the electron chemical potential upwards so that they cross in the hadronic freedom region between n_c and the flash chemical potential n_f . The dropping gauge coupling [19] and the rapid approach to

$a = 1$ in dense medium [45] enable one to reliably calculate the mass of the kaon as a function of density in the vicinity of the VM fixed point and hence determine the critical density. This phenomenon has a far-reaching consequence on the physics of compact stars. *What is important for our problem at hand is that when matter density is present, the vector dominance is more rapidly violated than in temperature alone.* This is highly relevant in our picture in NA60.

IV. DILEPTONS

We now apply the hadronic freedom picture to dileptons produced in relativistic heavy ion collisions. Our conclusion will be that if the hadronic freedom picture is correct, then the dileptons produced from the ρ mesons populating the interval between T_c and T_f are strongly suppressed and hence the measured dileptons do not provide a direct snapshot of the properties of the vector mesons that are BR-scaled.

A. The spectral function near the VM fixed point

Consider first the thermal ρ spectral function in HLS₁ theory in the close vicinity of the VM fixed point approached from below. We shall show that depending upon whether the ρ width is vanishing or non-vanishing, one gets a completely different result for the spectral function $\Re$.

Suppose that we have a renormalized ρ propagator in the form

$$D_\rho(q^2) = \frac{-1}{q^2 - m_\rho^{*2} + i\sqrt{q^2}\Gamma_\rho^*} \quad (28)$$

with

$$m_\rho^{*2} = M_\rho^{*2} + R. \quad (29)$$

where M_ρ^* is the mass parameter with intrinsic temperature dependence in the HLS₁ Lagrangian given by (14), R is the real part of the self-energy and Γ_ρ^* is the width coming from the imaginary part of the self-energy. The asterisk stands for an in-medium quantity. To make the discussion more general than HLS₁, we will suppose that the self-energy term could contain contributions from degrees of freedom other than just ρ and π . For instance, Γ_ρ^* could in principle contain many-body effects such as collisional width, involving baryonic excitations. The spectral function relevant for the dilepton spectrum at a fixed T (including $T > T_f$) is expressed by the imaginary part of the vector current correlator as

$$\Re \equiv \text{Im} G_V(q^2 = s) = \text{Im} [F_\rho^2(s) D_\rho(s)] \quad (30)$$

where $F_\rho(s)$ is the ρ decay constant which in general will have an imaginary part. Ignoring the imaginary part, we

have

$$F_\rho^2(s) = f_\sigma^{*2} \quad (31)$$

where f_σ^* is the in-medium decay constant of the longitudinal component of ρ (see Eq.(4)). Note that f_σ^{*2} includes both elementary hadronic and thermal loop corrections. It is related to the in-medium ρ - γ coupling g_ρ^* as

$$g_\rho^{*2} = m_\rho^{*2} f_\sigma^{*2}. \quad (32)$$

Setting $s = m_\rho^{*2}$ at the in-medium pole position, we get for the spectral function

$$\Re(s = m_\rho^{*2}) = \frac{g_\rho^{*2}}{m_\rho^{*3} \Gamma_\rho^*}. \quad (33)$$

In terms of the leptonic decay width which is given by

$$\Gamma_{ee}^* = \frac{4\pi\alpha^2}{3} \frac{g_\rho^{*2}}{m_\rho^{*3}}, \quad (34)$$

we have

$$\Re(s = m_\rho^{*2}) \sim \frac{\Gamma_{ee}^*}{\Gamma_\rho^*}, \quad (35)$$

where we have neglected certain numerical factors such as $\frac{4\pi\alpha^2}{3}$. This shows that the behavior of the spectral function depends on the properties of both the leptonic and hadronic widths.

Let us see what happens when one approaches the chiral transition point at which $g^* = f_\pi^* = f_\sigma^* = 0$. For this, we look at Eq. (33):

$$\Re(s = m_\rho^{*2}) \propto \frac{f_\sigma^{*2}}{g^* \Gamma_\rho^*} \quad (36)$$

where we have used $m_\rho^* \sim g^*$ near the fixed point. We need to know (a) how g^* and f_σ^* approach the fixed point and (b) more crucially the behavior near the fixed point of the hadronic width Γ_ρ^* . In HLS₁, it is established that $g^* \propto \langle \bar{q}q \rangle \rightarrow 0$ but how f_π^* (or f_σ^*) behaves in terms of the quark condensate is not known. It is because it goes to zero at the fixed point by the cancelation between the intrinsic term F_π^* and thermal loop corrections, both of which are not zero.

Suppose for the sake of argument that the in-medium pion decay constant goes as $f_\pi^* \sim g^*$. Then the behavior of the spectral function will be entirely dependent on whether the hadronic width vanishes or not. If it is non-vanishing, then the spectral function will go to zero. However if it vanishes, how it vanishes will matter. Within HLS₁, we expect $\Gamma_\rho^* \sim g^{*2}$. In this case the spectral function will be singular. This would be the case if leptons decoupled from the ρ after hadrons (e.g., pions) did. If this case holds, then the quantitative estimate made on the suppression factor discussed in section V would have to be taken with caution.

In reality, the situation is not at all clear. First of all, there is the threshold mass required by the lepton mass (as in NA60) which prevents the approach to the fixed point. Furthermore there can be several elements or mechanisms that are not encoded in HLS₁ at the perturbative level which can intervene to modify the picture. We have already mentioned, among other things, the possible role of the infinite tower implied by holographic dual QCD, the solitonic degrees of freedom (e.g., collisional broadening), Hagedorn excitations etc. This means that we cannot simply take the HLS₁ result mentioned above at its face value. Now given our ignorance of the variety of mechanisms that could possibly invalidate the simple HLS₁ prediction, we choose to accept the experimental indication, described in section V, that *no* dileptons are observed from the ρ mesons living in the temperature interval between T_c and T_f that we are identifying with hadronic freedom. It would of course be extremely interesting if future experiments did exhibit a sharp peak at a low dilepton invariant mass. It would provide an evidence for the validity of the simplest picture of HLS₁ and of the vector manifestation fixed point. It would however falsify the role of the assumed hadronic freedom on the strong suppression of the dilepton coupling to the vector mesons in the given temperature (and density) range.

B. Suppression of dileptons in the hadronic freedom

Although similar arguments apply to other dilepton experiments (such as CERES, PHENIX etc.), we will focus here on the conditions that are met in NA60.

There are several sources in NA60 for the dimuon suppression in the hadronic free region in temperature from T_c to T_f as well as in density from n_c to n_f . In this region, the ρ meson decay to a lepton pair is suppressed both in the EM sector and in the hadronic sector. Notable among the possible suppression mechanisms are:

1. **VD violation:** In the EM sector, the maximal violation of VD discussed above reduces the $\rho \rightarrow l^+l^-$ decay rate by a factor of 4. This reduction sets in more quickly in the presence of baryonic matter as mentioned in section IIIC.
2. **Wave function suppression:** At the VM fixed point, the transverse ρ decouples from the vector current as the gauge coupling g goes to zero. If we were to consider in terms of the quark picture, this would imply, in the hadronic sector, the suppression of the bound zero-mass quark-quark-antiquark wave function of the ρ meson, $|\Psi_\rho(0)|^2$. This effect is encoded in HLS₁ theory by the suppression of the $\gamma\rho$ coupling which gives a factor of ~ 2 reduction.
3. **Phase space suppression:** Because of the dropping gauge coupling g , $a \approx 1$ and the dimuon

threshold mass of 210 MeV, the phase space for the decay is suppressed by a factor ~ 2 . This effect is not entirely orthogonal to the wave function suppression effect as explained below.

These mechanisms will give a suppression factor relative to the standard picture [46] of the dileptons of ~ 10 in the hadronic freedom regime where the BR scaling effect is to be operative in HLS₁ theory.

We elaborate on these points.

1. VD violation

The violation of VD comes about by a^* going to 1 in hot/dense medium. Although one should perform the renormalization group calculation to find a^* in terms of temperature as well as density – such a calculation has not been done yet, there is a strong indication that a^* goes rapidly from 2 to 1 in hot/dense medium. One can see in the work of Shuryak and Brown [21] that this drop is sudden, at $T \approx 120$ MeV, with increasing T . In the presence of density, it should drop even faster as argued in section IIIC. Therefore, we assign $a^* = 1$ to the region from ~ 120 MeV to ~ 175 MeV. This gives the suppression factor of about 4. Moreover, there will be dileptons coming from the direct $\gamma\pi\pi$ coupling that have nothing to do with the in-medium properties of the ρ meson. These should contribute a roughly constant background to the total dilepton yield. Note that the range of ρ -meson energies is 0 to 700 MeV, so that we are discussing ρ -mesons with energy below the on-shell ρ -mass of 770 MeV. Therefore, we should look to the low mass part of the spectrum.

2. ρ wave function suppression

There will be further hindrance for the dileptons which come through the ρ meson on the hadronic side, because the dileptons are similar to a photon, to the extent that their masses are small compared with their invariant energy, so that there is an approximate factor

$$F = |\psi_\rho(0)|^2 \quad (37)$$

(where ψ_ρ is the ρ wavefunction) of the probability of the off-shell ρ meson being at its origin. This gives rise roughly to a factor of 2 reduction. This can be seen in HLS₁ as follows: The photon couples to the ρ with the coupling $\sim a^*g^*F_\pi^{*2}$ which goes to zero when the VM is approached, i.e., $g^* \rightarrow 0$. The vanishing of the photon coupling manifested in the vanishing gauge coupling represents the vanishing of the ρ wave function at the origin, $|\psi_\rho(0)|_{m_\rho^*=0} = 0$, when the mass goes to zero. What is relevant to us is the pion-loop medium correction δ to the $\gamma\rho$ coupling $g_{\gamma\rho}$

$$\frac{g_{\gamma\rho}^*}{g_{\gamma\rho}} \sim \frac{a^*g^*(1+\delta)}{ag} \quad (38)$$

with $F_\pi^*/F_\pi \approx 1$. The density dependence of δ has not yet been computed in HLS₁ but its temperature dependence has been computed perturbatively in Ref. [33]:

$$\begin{aligned}\delta(T = T_f) &\approx -0.2, \\ \delta(T = T_c) &\approx -0.4\end{aligned}\quad (39)$$

for $T_f \approx 120$ MeV and $T_c \approx 175$ MeV. This increases the reduction factor by an additional factor $\sim 1.6 - 2.8$. Given that the density effect is not included in the loop correction, it seems reasonable to take the resulting reduction factor to be ~ 2 .

We have to stress here that the estimates (39) were made in perturbation theory with HLS₁ Lagrangian with ρ and π only, so may not be reliable in the vicinity of the phase transition point. There may be non-perturbative effects which require going beyond the chiral perturbative approach to HLS theory. Furthermore, the drastically simplified HLS structure with the ground state vector mesons only may be inadequate in describing dynamics near the phase transition. As mentioned above, such degrees of freedom as the infinite tower of vector mesons encoded in holographic QCD, baryonic excitations via solitons (i.e., skyrmion or instanton) etc. might play a crucial role.

3. Phase space suppression

One can arrive at a similar reduction factor from the phase space consideration which seems to have a partial overlap with the wave function suppression considered above. Integrating the contribution from potentially BR-scaling ρ mesons down from T_c , we first find that the integral begins at $2m_\mu \sim 210$ MeV, because the muon masses must be furnished, and then continues down to the nearly on-shell ρ mass of ~ 700 MeV, the remaining 10% to put the ρ completely on shell coming from the kinetic energy [21]. However, in our scenario, the ρ masses are essentially zero [47] at just above T_c [26] behaving as if they have a quark substructure of two massless quarks, coupled to a massless ρ . In the integral going down from T_c the integrand, beginning from zero at $2m_\mu$, would be expected to increase as the ρ goes back towards the value of the on-shell ρ because of the effect on $\bar{\psi}_\rho\psi_\rho(0)$ from the increasing correlations as the ρ goes on shell. It seems reasonable to approximate the entire effect of $\bar{\psi}_\rho\psi_\rho(0)$ by a linearly increasing function which starts from zero at $2m_\mu$ and goes to 1 at $T = T_{\text{flash}} = 120$ MeV, where the ρ mesons go on shell. This will cut the dileptons that go through the ρ spectral function down by a factor of ~ 2 .

V. DILEPTONS “SEEN” IN THE EXPERIMENT

We have argued that the dileptons produced from the ρ mesons between $T_c \approx 175$ MeV and $T_f \approx 120$ MeV are strongly suppressed. The question then is: What are the

dileptons observed in NA60 (and also in CERES)? We suggest that the answer to this is that they are almost, if not entirely, *produced outside* of the hadronic freedom region, namely, at the flash temperature T_f . We present arguments to support this suggestion.

A. Dileptons in the vicinity of the $\rho(770)$

For our purpose, we consider most useful the article on radial flow of thermal dileptons by R. Arnaldi *et al.* [48]. In Fig.2 is reproduced their plot of the T_{eff} obtained in an effective theory along the line of flow, which produces a blue-shifting

$$T_{\text{eff}} = \sqrt{\frac{1+\beta}{1-\beta}} T \quad (40)$$

and in which the Boltzmann factor m is replaced by $m_\perp$, the mass perpendicular to the flow. Here β refers to the flow. Since the flow develops only late, after the hadrons have gone on shell, the amount of blue shift tells one about how late the hadrons emerge [49]. By disentan-




FIG. 2: Out of the blue-shifted hadrons, the dileptons from the ρ are shifted the most. They are reconstructed on the (steepest) dashed line. Between the dashed and solid lines, the ω , ϕ and η will flow somewhat less than the ρ 's, but nonetheless are blue shifted. The T_{eff} for the ρ has 300 ± 17 MeV for the peak, and 231 ± 7 MeV for the underlying continuum in the window $0.6 < M < 0.9$ MeV. Reproduced from Fig. 4 of [48].

gling the peak from the continuum Arnaldi *et al.* find $T_{\text{eff}} = 300 \pm 17$ MeV for the peak and 231 ± 7 MeV for the underlying continuum in the window $0.6 < M_{\mu\mu} < 0.9$ MeV. The latter is suggested as coming from the η , ω , and ϕ , which freeze out earlier due to their smaller coupling to the pions, the latter producing the flow. Note that in our scenario, the ω and η are included in the $SU(4)$ of particles which go massless at T_c .

The ρ -meson that has been reconstructed as the peak on the broad continuum [50] is shown in Fig. 3. We

can interpret this as the ρ meson that emerges from the hadronic free region and goes on-shell at the flash point. To confirm the validity of this interpretation, let the peak height at the on-shell mass $m_\rho = 770$ MeV – which fixes the normalization – be denoted by D which we take to be $D \approx 620 \times 10^3$ from Fig. 3. We take the quantity given by Fig. 3 to be of the form

$$\text{Fig. 3} = DF(M_{\mu\mu}) \left(\frac{m_\rho}{M_{\mu\mu}} \right)^2 \quad (41)$$

where $M_{\mu\mu}$ is the dimuon invariant mass and F is the relativistic form of the Breit-Wigner formula,

$$F(M_{\mu\mu}) = \left(1 + \left\{ \frac{1 - (M_{\mu\mu}/m_\rho)^2}{\Gamma_\rho/m_\rho} \right\}^2 \right)^{-1}. \quad (42)$$

Let us see whether Fig. 3 coincides with or deviates appreciably from the spectral function of the on-shell ρ with the free-space ρ width $\Gamma_\rho = 150$ MeV. This can be easily checked. Taking $M_{\mu\mu} = 400, 600$ and 780 MeV in Eq. 42, we find, in units of 10^3 , 153, 202 and 593 respectively. These are essentially what's given by Fig. 3, i.e., 160, 260 and 600. Thus modulo the normalization, the ρ^0 's in the flow are all on-shell ρ 's *unaffected* by the medium. There is evidence for neither BR-scaling ρ^0 's nor ρ^0 's with broadened widths.




FIG. 3: The on-shell ρ reconstructed by NA60 assuming $T_{\text{freezeout}} = 110$ MeV and $m_\perp$ scaling across the ρ resonance (not very sensitive). Reproduced from [50].

So far we have not considered explicitly what a_1 mesons can do. Dusling, Teany and Zahed [51] fit well the part of the dilepton spectrum below the ρ meson mass around $M = 500$ MeV by invoking effects from the a_1 meson at 1260 MeV. In our scenario, the a_1 meson makes up part of the 32 degrees of freedom, massless at T_c , that are flowing without interactions from T_c down to T_f . At T_f , the a_1 , having an immense on-shell width of ~ 200 -650 MeV, suddenly goes on shell, changing into

a $\rho + \pi$. This ρ joins the flow. This contribution which doubles the effective number of ρ^0 's at T_f is included in our scenario in the normalization of Fig. 3. The a_1 effect of Dusling et al. could however figure in our scenario outside of the hadronic freedom regime and outside of the flow as mentioned below.

B. Dileptons below the $\rho(770)$ peak

Let us now turn to how one can understand the dileptons that contribute to the spectral function *below* the free-space ρ mass. The principal thrust of this paper was that the bump near ~ 770 MeV is populated predominantly by near-on-shell ρ 's that are separated by the blue-shifting in the effective thermodynamics flow, and that in the invariant mass region below ~ 770 MeV in which BR scaling vector mesons should be present, the photon coupling is highly suppressed in accordance with the hadronic freedom. As a result, those dileptons produced by mundane mechanisms, i.e., “background,” should dominate the spectral function, masking more or less completely BR-scaled vector mesons. We suggest that the most likely mechanism that produces the dominant component of the “background” is pions in heat bath in strong interactions that populate at or just below the flash point with the Shuryak-Brown kinematical effects taken into account.

Our basic premise is that in the hadronic freedom scenario, all hadrons, including pions, flow freely from the critical temperature to the flash point in a time of $\gtrsim 3$ fm/c. (As mentioned, pions will also be weakly interacting like other hadrons, because the hidden gauge coupling g – which governs the p-wave $\pi\pi$ interaction through ρ exchange – is small.) In this regime, the photon has a direct coupling to a pion due to the violation of vector dominance with $a \sim 1$ – which would reflect the presence of the vector manifestation – but relatively few leptons are expected to be produced by the pions in that regime.

Reaching the flash point, however, all hadrons making up ~ 32 degrees of freedom going on-shell, with the hidden gauge coupling regaining its free-space strength, will produce dileptons with the vector dominance recovered with a going to 2. The vector mesons produced in the interactions at the flash point will go into flow and appear at on-shell mass.

Now in the central collisions, the freezeout temperature is lower than the flash temperature, so the hadrons which go on shell at the flash temperature go several generations before they freeze out at $T_{\text{freezeout}}$. The large number of pions present between T_f and $T_{\text{freezeout}}$ – not in the flow – will interact strongly producing, among others, copious ρ^0 's which will emit the lepton pairs. These ρ^0 's, interacting strongly in medium, will develop widths that are due to conventional nuclear interactions – including “sobars” [52] – in the form of the ρ self-energy. These will populate the invariant masses below the $\rho(770)$ peak. The a_1 's so produced could also emit dileptons à la

Dusling et al. We suggest that most, if not all, of what's calculated in the references [51, 53] belong to this class of contributions which have nothing to do with what's happening in the hadronic freedom regime.

If one considers the infinite-tower structure of HLS_∞ , there can be additional sources for dileptons of lower invariant mass. Let us consider how the higher members of the tower can contribute to the dilepton production. Let us first look at their contribution in the process $\pi^+\pi^- \rightarrow l^+l^-$ ($l = \mu, e$). The relevant graph is given by Fig. 4. This is given by the form factor (17) separated




FIG. 4: Dilepton ($l = e, \mu$) production by $\pi^+\pi^-$ annihilation in heavy ion collisions involving the infinite tower of vector mesons ρ_n $n = 0, 1, 2, \dots, \infty$

into two components as (22). Here we are dealing with the time-like regime for which $1/N_c$ corrections need to be included. Let us imagine that they can be calculated. Then they will give both real and imaginary parts. The real part will shift the masses and the imaginary part will give the widths. We expect that the low-mass component (23) will essentially give the Breit-Wigner form

$$F(M_{l^+l^-}^2) \approx \left(1 + \left\{ \frac{1 - (M_{\mu^+\mu^-}/m_{\rho(770)})^2}{\Gamma_\rho/m_{\rho(770)}} \right\}^2 \right)^{-1} \quad (43)$$

with the free-space ρ width $\Gamma_\rho = 150$ MeV. As for F^{high} , we can ignore the self-energy correction and $|q^2| = M_{\mu^+\mu^-}$ which are expected to be much smaller than the higher-tower vector-meson masses in (24) and find that due to the charge sum rule, Eq. (25) applies to the form factor in the time-like regime. It may be possible to justify this argument with the time-like form factor obtained from the τ decay measurement mentioned above [32]. Thus essentially the high tower does not affect the structure that is obtained in HLS_1 with $a = 2$ in the time-like regime as in the space-like regime.

What we have shown above is that the 2π contributions to dileptons are constrained by the pion charge sum rule involving the infinite tower of vector mesons in hQCD. However there are no such constraints for multi-pion contributions. Specifically in HLS_∞ , $2n\pi$ for $n > 1$ can contribute importantly via higher ρ_n 's. Of course there is the threshold effect so that the 4π contribution will be lower-bound by ~ 560 MeV. In fact there is indication that $4\pi \rightarrow e^+e^-$ receives substantial contributions from $\rho(1450)$ and $\rho(1700)$. This indication comes from a recent analysis of the available CMD-2 and BaBar data implementing the three lowest members of the tower and a_1

mesons [54]. The complete vector dominance structure of hQCD is not present in the analysis of [54], so it is not obvious what the infinite tower structure will do in heavy ion processes. We expect that such four-pion contributions mediated by higher members, if present, could make non-negligible contributions to dileptons with invariant masses $\gtrsim 560$ MeV.

VI. COMMENTS AND CONCLUSIONS

Brown-Rho (BR) scaling was a prediction based on effective field theory, not a hypothesis or a conjecture as often stated, in a phenomenological attempt to build the main feature of chiral invariance, the behavior of the broken chiral symmetry which gives hadrons, other than the pion, masses. A great deal of activity had been spent in the years preceding BR scaling to introduce quarks as substructure of the hadrons into nuclear physics. Whereas there were some successes, such as in the chiral bag model, color transparency, etc., there was not a broad theoretical basis for the many-body problem. A local field theory that did just that by matching to QCD was introduced by Harada and Yamawaki in HLS_1 which gave the first effective field theory support for BR scaling. BR scaling also came out naturally in strong-coupling lattice gauge calculations in the large $1/d$ expansion (where d is the number of space dimension) [55].

The original idea of BR scaling was that in medium, because of background density or temperature which affects the “vacuum” structure, hadron masses, other than that of the pion, that are dynamically generated had to be scaled as a function of the quark condensate. In many-body systems where such a vacuum change is generated by strong nuclear interactions, this idea can be addressed only in an effective field theory framework and not directly in QCD language. So the question then is: In what way would BR scaling manifest in complex many-body processes? The answer to this question depends on scales that are involved. This point was recognized in the early phenomenological papers by Brown and Rho where the scaling effect was looked for in spin-isospin dependent nuclear interactions, notably the tensor forces. Thus very near the chiral transition point, HLS_1 theory, well-defined by the VM fixed point, makes an extremely simple and clear-cut prediction – whether right or wrong – in the property of the vector mesons ρ and ω and hence of physical observables associated with them. However away from the VM fixed point, the situation is quite different and a lot more intricate. While the parameters of the HLS_1 Lagrangian are directly linked to the quark condensate thanks both to the local gauge invariance and to the matching to QCD, most physical observables, while dependent on BR scaling, have complicated dependence on the chiral order parameter. For instance, at the scale of the structure of light nuclei, BR scaling can figure in giving effective nuclear forces. Most striking of all in

this category is the archaeologically long lifetime of the ^{14}C decay explained in terms of the tensor force that results from an intricate cancelation between the pionic term and the ρ term where BR scaling enters through the parametric mass term [56]. On the other hand, near the Fermi liquid fixed point of nuclear matter, BR scaling manifests itself in the Landau Fermi liquid parameter F_1 , explaining how exchange currents affect the orbital gyromagnetic ratio in heavy nuclei [57]. It is in this context (as stressed in [52]) that one would have to understand the role of dileptons as a possible litmus indicator for chiral symmetry in hot and/or dense matter.

Using the notion of the hadronic freedom, we have argued that the dileptons measured in heavy ion collisions are mostly, if not all, from the ρ^0 's outside of the hadronic freedom (or fireball) regime. We should stress that the hadronic freedom, a notion derived from HLS₁ with its VM fixed point, has an additional assumption extraneous to HLS₁ in the VM that requires verification. It assumes that the ρ width at the VM fixed point is not zero whereas a strict adherence to HLS₁ with no additional ingredients – which we believe is not complete – would imply that the width vanishes. This renders it to a possible falsification. An observation of a strong enhancement of the dilepton production in the *close vicinity* of T_c , while giving a partial support to the simplest HLS₁, would falsify the notion of the hadronic freedom and hence our scenario on dileptons described in this paper.

The violation of vector dominance, a characteristic of HLS₁ in the vector manifestation, could also be subject to falsification. While dileptons from ρ mesons are suppressed, one expects dileptons coming from pions via direct coupling due to the VD violation. To see this effect, it would be necessary to subtract *all* the “background” coming from on-shell sources. This might be a daunting task but if such effect is found, it will be a “smoking-gun” signal for HLS₁ in the vector manifestation and also an indirect support for BR scaling.

There have been discussions, both experimental and theoretical, on the behavior of vector mesons in cold dense nuclear medium [58]. Experiments in the upcoming accelerators will surely generate a lot of activities in this area. It would be difficult to probe directly the density regime in the close vicinity of the VM fixed point, so what one can do is to study various precursor effects of matter at densities near that of nuclear matter. The JLab [60], CD-Bonn [61] and KEK [62] experiments belong to this class.

We now check what is involved in HLS₁ theory for such effects.

What is addressed in this class of experiments is the “mass” of the vector mesons ρ and ω “seen” inside nuclear medium at a density $\lesssim n_0$. The measured quantities reflect the pole mass which in HLS₁ is given by what is equivalent to Eq. (15) in density,

$$m_V^{*2} = a^* F_\pi^{*2} g^{*2} + \Sigma^* \quad (44)$$

where the asterisk here stands for the *intrinsic* depen-

dence on the chiral order parameter, i.e., the quark condensate, which changes with density and Σ^* is the “dense loop” corrections gotten from the HLS₁ Lagrangian in a suitable perturbative series. Now to address the dense loop corrections, it is necessary to introduce fermionic degrees of freedom to implement matter density. In principle this could be done in terms of dense skyrmions in HLS₁ or better in HLS_∞ involving the infinite tower of vector mesons in 4D or instantons in 5D which automatically takes into account the intricate vacuum structure in which both pions and vector mesons are condensed generating a self-consistent baryonic background for fluctuating mesons. In a sophisticated version of this type of approach, one could also build in N^* 's, thereby incorporating such collective excitations as “sobars” [52] etc. Efforts are being made to arrive at dense matter through this approach but up to date there has been little progress along this line.

What has been done up to date is to introduce, somewhat ad hoc, quasiquarks in a gauge invariant way into HLS₁ [19]. At one loop order, we can rewrite (44) as

$$m_V^{*2} = a^* f_\pi^{*2} g^{*2} + 12g^{*2} c G^* \quad (45)$$

where f_π^* is the pion decay constant renormalized by dense one-loop corrections, G^* is a known function of density and c is the only undetermined constant in the Lagrangian which is of $\mathcal{O}(1)$. We can make a rough estimate of (45) at nuclear matter density [63],

$$\begin{aligned} (m_V^*/m_V)^2(n_0) &\approx (f_\pi^*(n_0)/f_\pi)^2 + 0.21c \\ &\approx 0.64 + 0.21c \end{aligned} \quad (46)$$

where we have used the value $f_\pi^*/f_\pi \approx 0.8$ quoted in the literature at nuclear matter density [64]. If $c > 0$ and $c \sim \mathcal{O}(1)$, one sees that the dense loop corrections – that comprise essentially of mundane nuclear interactions in the given field theory framework – are comparable to the first term that purports to provide direct information on BR scaling. We see from this simple calculation that in order to reveal unequivocally the role of BR scaling, one first has to define precisely what quantities one is to zero-in on in a precisely defined effective field theory, and then to compute many-body corrections in a way consistent with the effective theory so defined. Phenomenological Lagrangians constructed solely to fit free-space quantities – hence valid at tree order and devoid of the Wilsonian renormalization-group flow [5] – are not likely, if not impossible, to provide useful information of this subtle nature. In HLS₁, it is the parametric mass (first term on the RHS of (45)) – and not the measured pole mass – that carries the looked-for information.

That dileptons, considered to be an ideal snapshot of the chiral order parameter, appear to be blind to BR scaling may seem to indicate that BR scaling cannot be probed unambiguously. This is however not the case. There are in fact certain quantities that could provide a clear-cut signal for the BR scaling at work. One such case that has been studied is the pion velocity v_π at T_c . The ρ

mass becoming degenerate with the pion mass as $T \rightarrow T_c$ in HLS₁ in the VM makes the pion velocity rapidly approach 1 [65], whereas in the absence of the massless ρ , v_π can be rigorously shown to vanish [66]. This is a night-and-day difference. The pion velocity going to zero would be a strong indication that the ρ mass does not drop, hence the vector manifestation is not realized in nature and would call for a revamping of the hidden gauge structure in the form of HLS₁, perhaps requiring the infinite tower of vector mesons. It would be a challenge to both theorists and experimentalist to find other cases where HLS₁ in the vector manifestation can be exposed equally directly and ultimately to exploit HLS_∞ in exploring the intricate nature of matter near the critical point. In this connection, it is of importance to note that in holographic QCD, baryons emerge as coherent states of pions and vector mesons, in particular, the ρ meson, and the vector modes “seen” in medium are excitations on top of this ground state in which both pions and vector mesons are condensed. Thus the property of vector mesons in baryonic medium is expected to be vastly more

intricate than the naive picture would give.

Acknowledgments

Three of us (MH, MR and CS) acknowledge the hospitality of Yukawa Institute of Theoretical Physics where “Yukawa International Program for Quark-Hadron Sciences” entitled “*New Frontiers in QCD 2008*” was held and where a part of this paper was written. The work of MH was supported in part by the JSPS Grant-in-Aid for Scientific Research (c) 20540262 and the Global COE Program “Quest for Fundamental Principles in the Universe” of Nagoya University provided by Japan Society for the Promotion of Science (G07), part of the work of MR was funded by the WCU project of the Korean Ministry of Educational Science and Technology and CS acknowledges partial support by DFG cluster of excellence “Origin and Structure of the Universe”.

-
- [1] G.E. Brown and M. Rho, Phys. Rev. Lett. **66** (1991) 2720.
 - [2] See, e.g., A. Ferretti et al., *Proceedings of 42nd Rencontres de Moriond on QCD and Hadronic Interactions*, La Thuile, Italy, 17-24 March 2007, e-Print: arXiv:0705.3339 [nucl-ex] where all previous references to the experiment are found.
 - [3] By “mundane” nuclear processes, we mean those processes involving hadrons, both mesons and baryons, in nuclear medium that are calculable starting with a phenomenological chiral Lagrangian – possessing pertinent symmetries – whose parameters are fixed in the matter-free vacuum. By definition, calculations are strictly valid at tree order although in practice loops are commonly computed with phenomenological form factors or cut-offs.
 - [4] Unless otherwise stated, a similar argument applies to density although we will address only temperature.
 - [5] M. Harada and K. Yamawaki, Phys. Rept. **381** (2003) 1.
 - [6] H. Leutwyler, “On the low energy end of the QCD spectrum,” arXiv:0809.5053 [hep-ph].
 - [7] In nature, the quark mass is not zero, so we are not at the chiral limit but barring some special cases, it is a good approximation. We will be considering the chiral limit unless otherwise stated.
 - [8] Going systematically to higher energy scales with gauge-fixed Lagrangians is known to be very difficult if not impossible. It is the local gauge invariance that allows a systematic account of higher derivative terms and renders the effective theory amenable to a UV-completion.
 - [9] N. Arkani-Hamed, H. Georgi and M.D. Schwartz, Annals Phys. **305** (2003) 96.
 - [10] Here N_f is for any number of flavors but will be taken to be $N_f = 2$ for applications.
 - [11] M. Bando, T. Kugo, S. Uehara, K. Yamawaki and T. Yanagida, Phys. Rev. Lett. **54** (1985) 1215.
 - [12] M. Bando, T. Kugo and K. Yamawaki, Phys. Rept. **164** (1988) 217.
 - [13] D.T. Son and M. Stephanov, Phys. Rev. **D69** (2004) 065020.
 - [14] T. Sakai and S. Sugimoto, Prog. Theor. Phys. **113** (2005) 843; Prog. Theor. Phys. **114** (2005) 1083.
 - [15] M. Harada, S. Matsuzaki and K. Yamawaki, Phys. Rev. **D74** (2006) 076004.
 - [16] Going beyond one-loop order is highly daunting and at present is unpracticable not only because there are a large number of diagrams that enter but also a large number of parameters to be fixed, which is not feasible given the present state of experimental data. It should however be mentioned that the results given below are valid at least qualitatively to all orders of chiral expansion.
 - [17] S. Weinberg, Phys. Rev. **D56** (1997) 2303.
 - [18] M. Harada and C. Sasaki, Phys. Lett. **B537** (2002) 280.
 - [19] M. Harada, Y. Kim and M. Rho, Phys. Rev. **D66** (2002) 016003.
 - [20] The parameteric mass will be denoted as M_ρ while the physical (pole) mass will be given as m_ρ .
 - [21] E.V. Shuryak and G.E. Brown, Nucl. Phys. **A717** (2003) 322.
 - [22] P. Fachini, R. S. Longacre, Z. Xu, and H. Zhang, J. Phys. G: Nucl. Part. Phys. **34** (2007) 431.
 - [23] D.E. Miller, Phys. Rept. **443** (2007) 55.
 - [24] It will be more direct to determine the flash temperture from lattice measurements of the quark condensate. We became aware of the measurement of the light-quark condensate for reasonably realistic quark masses, M. Cheng *et al*, Phys. Rev. **D77** (2008) 014511, after the preliminary version of this paper was posted on the web. This measurement would give a flash temperature ~ 140 MeV. We would like to thank Wolfram Weise for pointing out this reference to us.
 - [25] G.E. Brown, C.-H. Lee, M. Rho, “The STAR ρ^0/π^- ratio in Au-Au peripheral collisions at RHIC and the vector manifestation of hidden local symmetry,” nucl-th/0507011.

- [26] H.-J. Park, C.-H. Lee and G.E. Brown, Nucl. Phys. **A763** (2005) 197.
- [27] An interesting possibility to explore is that the existence of the infinite tower of mesons in the hadronic freedom regime could figure in cooling of the system from T_c to T_f .
- [28] J.J. Sakurai, Currents and mesons (Chicago U.P., Chicago, 1969).
- [29] M. Harada and K. Yamawaki, Phys. Rev. Lett. **87** (2001) 152001 [arXiv:hep-ph/0105335].
- [30] M. Harada and C. Sasaki, Nucl. Phys. **A736** (2004) 300.
- [31] D.K. Hong, M. Rho, H.-U. Yee and P. Yi, JHEP **09** (2007) 063; Phys. Rev. **D77** (2008) 014030; M. Rho, “Baryons and vector dominance in holographic dual QCD,” arXiv:0805.3342 [hep-ph]; J. Park and P. Yi, JHEP **0806** (2008) 011; P. Yi, “Holographic baryons,” to appear.
- [32] T.E. Coen et al, Phys. Rev. Lett. **92** (2004) 232001. In this article, the process of τ decay into $K^-\pi^-K^+\nu_\tau$ was studied. From Eq. (7) of this article, we can extract the pion charge form factor in the form (in the notation of this reference)
- $$F(Q^2) = \frac{BW_\rho(Q^2) + \lambda BW_{\rho'}(Q^2) + \delta BW_{\rho''}(Q^2)}{1 + \lambda + \delta}$$
- where $\rho = \rho_0$, $\rho' = \rho_1$ and $\rho'' = \rho_2$ in our notation. Best fit to experiments for $1.2 \text{ GeV} < \sqrt{Q^2} < 1.8 \text{ GeV}$ gave $\lambda = -0.314 \pm 0.073$ and $\delta = 0.101 \pm 0.020$. Assuming that the momentum dependence can be ignored, this would give for the charge sum rule (19)
- $$F(0) = 1.27 - 0.40 + 0.13 + 0.00 = 1.00. \quad (47)$$
- Although one might suspect that there could be appreciable momentum dependence, particularly in the time-like regime where the constants λ and δ are determined, this is tantalizingly close to the Sakai-Sugimoto result (21). It also provides a support to the argument developed later for pion-lepton couplings. It would be interesting to take into account the momentum dependence and check whether this result survives.
- [33] M. Harada and C. Sasaki, Phys. Rev. **D74** (2006) 114006.
- [34] STAR Collaboration, Phys. Rev. Lett. **92** (2004) 092301.
- [35] R. Rapp, unpublished.
- [36] P. Braun-Munzinger, K. Redlich and J. Stachel, in *Quark Gluon Plasma 3* eds. R. C. Hwa and X.-N. Wang (World Scientific Publishing), nucl-th/0304013.
- [37] G.E. Brown, C.-H. Lee and M. Rho, Phys. Rev. **C74** (2006) 024906.
- [38] M. Asakawa, T. Hatsuda and Y. Nakahara, Nucl. Phys. **A715** (2003) 8630.
- [39] As explained in detail in [37], we are taking into account scalar and axial-vector degrees of freedom in the HLS framework although their inclusion has not been rigorously formulated.
- [40] S. Weinberg, Phys. Rev. Lett. **65** (1990) 1177; “Unbreaking symmetries,” in *Salamfestschrift* ed. A. Ali, J. Ellis and S. Randjbar-Daemi (World Scientific, Singapore, 1994).
- [41] This suggests yet another feature that departs drastically from the standard scenario that depicted a strongly-correlated hadronic matter below T_c making transition to a non-interacting quark gluon plasma above T_c . Our scenario instead has it that a non-interacting nearly massless hadronic matter below T_c makes a transition to a strongly-correlated matter just above T_c .
- [42] G.E. Brown, C.-H. Lee and M. Rho, Nucl. Phys. **A747** (2005) 530.
- [43] G.E. Brown, B. Gelman, M. Rho, Phys. Rev. Lett. **96** (2006) 132301.
- [44] G. E. Brown, C. H. Lee and M. Rho, Phys. Rev. Lett. **101** (2008) 091101 [Phys. Rev. Lett. **101** (2008) 119901]; Phys. Rept. **462** (2008) 1.
- [45] M. Rho, “Hidden local symmetry and dense half-skyrmion matter,” e-Print: arXiv:0711.3895 [nucl-th].
- [46] By “standard picture” we mean the description in which vector dominance is assumed and mundane many-body effects are taken into account in both the real and imaginary part of self-energy of the vector meson in the temperature interval involved. It may or may not incorporate some sort of “dropping mass” effects. As noted above, this factor cannot be applied without suitable modifications to the HLS₁ result of [33].
- [47] The $SU(4)$ symmetry at just above T_c is slightly broken by the spin-dependent force, but the inertial parameter p_0 is large, so the mass is only a few percent of the free ρ mass.
- [48] R. Arnaldi *et al.*, Phys. Rev. Lett. **100** (2008) 022302.
- [49] We are not concerned with the sudden decline (and therefore flow) in T_{eff} at masses $> 1 \text{ GeV}$ which is most naturally explained as a transition to partonic processes like $\bar{q}q \rightarrow \mu^+\mu^-$.
- [50] S. Damjanovic, in http://www.ect.it/Meetings/ConfWksAndCollMeetings/ConfWksDocument/2007/Talks/18_22_June/Damjanovic.cut.ppt.
- [51] K. Dusling, D. Teany and I. Zahed, Phys. Rev. C **75** (2007) 024908.
- [52] G.E. Brown and M. Rho, “NA60 and BR scaling in terms of the vector manifestation: Formal consideration,” e-Print: nucl-th/0509002; “NA60 and BR scaling in terms of the vector manifestation: A Model approach,” e-Print: nucl-th/0509001.
- [53] H. van Hees and R. Rapp, “Medium modifications of vector mesons and NA60,” arXiv:hep-ph/0604269; J. Ruppert and T. Renk, Eur. Phys. J. C **49**, 219 (2007); J. Ruppert, C. Gale, T. Renk, P. Lichard and J. I. Kapusta, Phys. Rev. Lett. **100** (2008) 162301.
- [54] N.N. Achasov and A.A. Kozhevnikov, “Hidden local symmetry and the reaction $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$ at energies $\sqrt{s} \leq 1 \text{ GeV}$,” arXiv:0809.3883.
- [55] A. Ohnishi, N. Kawamoto and K. Miura, “Brown-Rho scaling in the strong coupling lattice QCD,” arXiv:0803.0255[nucl-th].
- [56] J.W. Holt, G.E. Brown, T.T.S. Kuo, J.D. Holt and R. Machleidt, Phys. Rev. Lett. **100** (2008) 062501.
- [57] B. Friman and M. Rho, Nucl. Phys. **A606** (1996) 303.
- [58] Finite 3-momenta carried by ρ mesons are expected to affect in-medium ρ spectral functions and hence obscure the manifestation of chiral symmetry in nuclear medium. The calculation performed in hadronic many-body theory for the ρ produced by photon reactions off nuclei [59] has indeed shown that the rela-

tively large 3-momenta relevant for the JLAB CLAS experiments significantly affect the pole structure of the propagator in medium and hence the spectral function of the ρ meson. Thus, the dilepton rates with a low momentum are more relevant to details of medium effects in the data.

- [59] F. Riek, R. Rapp, T. S. Lee and Y. Oh, arXiv:0812.0987 [nucl-th].
- [60] R. Nasseripour *et al.* [CLAS Collaboration], Phys. Rev. Lett. **99** (2007) 262302.
- [61] D. Trnka *et al.* [CBELSA/TAPS Collaboration], Phys. Rev. Lett. **94** (2005) 192303.
- [62] K. Ozawa *et al.* [E325 Collaboration], Phys. Rev. Lett. **86**, 5019 (2001); M. Naruki *et al.*, Phys. Rev. Lett. **96**, 092301 (2006).
- [63] Let us rewrite Eq. (46) in a form that would be given in the standard (phenomenological) approaches with no local gauge symmetry and without matching to QCD – hence no intrinsic density dependence – as $(m_V^*/m_V)^2 = 1 + \Delta$. It is the Δ that the standard approaches would interpret – and the experimentalists measure – as the “mass shift.” In our simple calculation with HLS_1 , it is given by $\Delta = 0.12c - 0.36$. We see that for $c > 0$ and $c \sim \mathcal{O}(1)$, the “mass shift” can be negligible due to cancelation, the extent to which the cancelation takes place depending upon the detailed structure of the nuclei studied. In this form, BR scaling is not “visible.” This illustrates the difficulty in probing BR scaling at a density far from the VM fixed point.
- [64] P. Kienle and T. Yamazaki, Prog. Part. Nucl. Phys. **52** (2004) 85. In the Lorenz non-invariant system we are dealing with, the pion decay constant that figures is the time component. The difference between the time and space components of the pion decay constant – which is model-dependent as one can see in the pion velocity [65, 66] – shows up at higher orders than what we are working with, so we shall ignore it here.
- [65] M. Harada, Y. Kim, M. Rho and C. Sasaki, Nucl. Phys. **A727** (2003) 437.
- [66] D. T. Son and M. A. Stephanov, Phys. Rev. Lett. **88** (2002) 202302; Phys. Rev. D **66** (2002) 076011.